

PREVENTIVE CARE SERVICES GUIDELINES

SHPO reimburses providers for services rendered below as preventive benefits with a \$0.00 cost share to the member. If coded as indicated in the model, members pay \$0.00 for the service. Providers may also append Modifier 33 to indicate a preventive service as appropriate in accordance with billing standards.

SERVICE & DESCRIPTION	USPSTF RECOMMENDATION	CPT/HCPCS CODES	ICD-9-CM CODES	FREQUENCY & CRITERIA
Annual Wellness Exam – Assessment and Laboratory Services	N/A	99201-99215, 99381 – 99429, or G0402, G0438, G0439, S0610 - S0613 appropriate to the service provided and lab tests ordered or conducted during the visit	V70.0, V72.62, V20.2, V72.31, V76.2	All members. <i>1 time per benefit period</i>
Abdominal Aortic Aneurysm Screening: Men	The USPSTF recommends one-time screening for abdominal aortic aneurysm by ultrasonography in men aged 65 to 75 who have ever smoked.	76700, 76705, 76770, 76775, G0389	V15.82	Males age 65-75 who have ever smoked. <i>1 time per benefit year</i>
Alcohol Misuse Counseling	The USPSTF recommends screening and behavioral counseling interventions to reduce alcohol misuse by adults, including pregnant women, in primary care settings.	G0442-G0443 or service is included in a preventive care visit. Codes: 99381-99429 appropriate to the service	None	Men and Women age 18+. <i>1 time per benefit year</i>
Anemia Screening: Pregnant Women	The USPSTF recommends routine screening for iron deficiency anemia in asymptomatic pregnant women.	85004 -85041	Payable with diagnosis of pregnancy	Pregnant Women. <i>1 time per benefit year</i>

PREVENTIVE CARE SERVICES GUIDELINES (continued)

SERVICE & DESCRIPTION	USPSTF RECOMMENDATION	CPT/HCPCS CODES	ICD-9-CM CODES	FREQUENCY & CRITERIA
Aspirin To Prevent CVD: Men and Women	The USPSTF recommends the use of aspirin for men age 45 to 79 years and women age 55 to 79 years when the potential benefit due to a reduction in myocardial infarctions outweighs the potential harm due to an increase in gastrointestinal hemorrhage.	Service is included in a preventive care visit. Codes: 99381-99429 appropriate to the service	None	Men, Ages 45-79 and Women, Ages 55-79. <i>1 time per benefit year</i>
Bacteriuria Screening: Pregnant Women	The USPSTF recommends screening for asymptomatic bacteriuria with urine culture for pregnant women at 12 to 16 weeks' gestation or at the first prenatal visit, if later.	87081 - 87088	Payable with diagnosis of pregnancy	Pregnant Women at 12 to 16 weeks' gestation or at the first prenatal visit, if later. <i>1 time per benefit period</i>
Blood Pressure Screening	The USPSTF recommends screening for high blood pressure in adults aged 18 and older.	Service is included in a preventive care visit. Codes: 99381-99429 appropriate to the service	None	Men and Women age 18+. <i>1 time per benefit period</i>
Breast Cancer Genetic Risk Assessment and BRCA Mutation Testing (Counseling)	The USPSTF recommends that women whose family history is associated with an increased risk for deleterious mutations in BRCA1 or BRCA2 genes be referred for genetic counseling.	Service is included in a preventive care visit. Codes: 99381-99429 appropriate to the service or 96040 and S0265	V16.3 or V16.41	Women with a family history of ovarian cancer or breast cancer. <i>1 time per benefit period</i>

PREVENTIVE CARE SERVICES GUIDELINES (continued)

SERVICE & DESCRIPTION	USPSTF RECOMMENDATION	CPT/HCPCS CODES	ICD-9-CM CODES	FREQUENCY & CRITERIA
Breast Cancer Genetic Risk Assessment and BRCA Mutation Testing (Testing)	The USPSTF recommends that women whose family history is associated with an increased risk for deleterious mutations in BRCA1 or BRCA2 genes be referred for evaluation for BRCA testing.	S3818 - S3823, 83890 - 83913, 88271 - 88273, 88363 - 88365	V16.3 or V16.41	Women with a family history of ovarian cancer or breast cancer. <i>1 time per benefit period</i>
Breast Cancer Preventive Medication	The USPSTF recommends that clinicians discuss chemoprevention with women at high risk for breast cancer and at low risk for adverse effects of chemoprevention. Clinicians should inform patients of the potential benefits and harms of chemoprevention.	Service is included in a preventive care visit. Codes: 99381-99429 appropriate to the service	None	Women at high risk for breast cancer and at low risk for adverse effects of chemoprevention. <i>1 time per benefit period</i>
Breast Cancer Screening	The USPSTF recommends screening mammography for women, with or without clinical breast examination, every 1-2 years for women aged 40 and older.	77055, 77056, 77057, 77051, 77052, G0202, G0204, G0206	None	Women age 40+. <i>1 time per benefit period</i>
Breastfeeding Counseling	The USPSTF recommends interventions during pregnancy and after birth to promote and support breastfeeding.	Service is included in a preventive care visit. Codes: 99381-99429 appropriate to the service	None	Pregnant Women or Postpartum Women. <i>3 times per benefit period</i>
Cervical Cancer Screening	The USPSTF strongly recommends screening for cervical cancer in women who have been sexually active and have a cervix.	G0101, G0123 - G0124, G0141- G0148, P3000 - P3001, Q0091, 88141 - 88143, 88147 - 88155, 88164 - 88167, 88174 - 88175		Women age 18+. <i>1 time per benefit period</i>

PREVENTIVE CARE SERVICES GUIDELINES (continued)

SERVICE & DESCRIPTION	USPSTF RECOMMENDATION	CPT/HCPCS CODES	ICD-9-CM CODES	FREQUENCY & CRITERIA
Chlamydial Infection Screening: Women	The USPSTF recommends screening for chlamydial infection for all sexually active pregnant and non-pregnant young women aged 24 and younger and for older pregnant and non-pregnant women who are at increased risk.	86631 - 86632, 87110, 87270, 87320, 87490 - 87492, 87810		Women age 18+. <i>1 time per benefit period</i>
Cholesterol Abnormalities Screening: Men and Women	The USPSTF recommends screening men aged 20 to 35 and 35 and older for lipid disorders if they are at increased risk for coronary heart disease. The USPSTF also recommends screening women aged 20 to 45 and 45 and older for lipid disorders if they are at increased risk for coronary heart disease.	80061, 82465, 83718 -83721, 84478	V17.3, V17.49, V77.91, V81.0, V81.1, V81.2	Men, Ages 20 + and Women, Ages 20 +. <i>1 time per benefit period</i>
Colorectal Cancer Screening	The USPSTF recommends screening for colorectal cancer using fecal occult blood testing, sigmoidoscopy, or colonoscopy, in adults, beginning at age 50 years and continuing until age 75 years. The risks and benefits of these screening methods vary.	G0104 - G0106, G0120 - G0122, G0328, 44388 - 44394, 45330 - 45339, 45355 - 45385, 82270, 82274, 88304 - 88305	V76.41 or V76.51	Men and Women ages 50 – 75 years. <i>1 time per benefit period as clinically appropriate based on clinical care guidelines</i>
Dental Caries Chemoprevention: Preschool Children	The USPSTF recommends that primary care clinicians prescribe oral fluoride supplementation at currently recommended doses to preschool children older than 6 months of age whose primary water source is deficient in fluoride.	Service is included in a preventive care visit. Codes: 99381-99429 appropriate to the service	None	Children ages 6 months to 5 years. <i>1 time per benefit period</i>

PREVENTIVE CARE SERVICES GUIDELINES (continued)

SERVICE & DESCRIPTION	USPSTF RECOMMENDATION	CPT/HCPCS CODES	ICD-9-CM CODES	FREQUENCY & CRITERIA
Depression Screening: Adolescents and Adults	The USPSTF recommends screening of adolescents (12-18 years of age) for major depressive disorder when systems are in place to ensure accurate diagnosis, psychotherapy (cognitive-behavioral or interpersonal), and follow-up. The USPSTF also recommends screening adults for depression when staff-assisted depression care supports are in place to assure accurate diagnosis, effective treatment, and follow-up.	Service is included in a preventive care visit. Codes: 99381-99429 appropriate to the service	None	Adolescents and Adults, ages 12 – 18 and 18+. <i>1 time per benefit period</i>
Diabetes Screening	The USPSTF recommends screening for type 2 diabetes in asymptomatic adults with sustained blood pressure (either treated or untreated) greater than 135/80 mm Hg.	82947 - 82952, 83036	V77.1 and/or payable with a diagnosis of high blood pressure	Men and Women ages 18+. <i>1 time per benefit period</i>
Folic Acid Supplementation	The USPSTF recommends that all women planning or capable of pregnancy take a daily supplement containing 0.4 to 0.8 mg (400 to 800 µg) of folic acid.	Service is included in a preventive care visit. Codes: 99381-99429 appropriate to the service	None	Women ages 18+. <i>1 time per benefit period</i>
Gonorrhea Prophylactic Medications: Newborn	The USPSTF strongly recommends prophylactic ocular topical medication for all newborns against gonococcal ophthalmia neonatorum.	Service is included in a preventive care visit. Codes: 99381-99429 appropriate to the service	None	Newborns age 0-3 months. <i>1 time per benefit period</i>

PREVENTIVE CARE SERVICES GUIDELINES (continued)

SERVICE & DESCRIPTION	USPSTF RECOMMENDATION	CPT/HCPCS CODES	ICD-9-CM CODES	FREQUENCY & CRITERIA
Gonorrhea Screening: Women	The USPSTF recommends that clinicians screen all sexually active women, including those who are pregnant, for gonorrhea infection if they are at increased risk for infection (that is, if they are young or have other individual or population risk factors).	87590 - 87592, 87850, 87800, 87081, 87210, 87070, 87077	Payable with diagnosis of pregnancy or V69.2, V74.5, V76.2, or V01.6	Women ages 18+ who have been sexually active, and/or are pregnant. <i>1 time per benefit period</i>
Healthy Diet Counseling	The USPSTF recommends intensive behavioral dietary counseling for adult patients with hyperlipidemia and other known risk factors for cardiovascular and diet-related chronic disease. Intensive counseling can be delivered by primary care clinicians or by referral to other specialists, such as nutritionists or dietitians.	97802 - 97804, S9470, G0270 - G0271, G0446 – G0447, 98960 or may be included in a preventive care visit. Codes: 99381-99429 appropriate to the service	Payable with a diagnosis of hyperlipidemia, and/or with a diagnosis related to cardiovascular disease, and/or a diet-related chronic disease	Men and Women age 18+ who have hyperlipidemia or other known risk factors for cardiovascular disease or diet-related chronic disease. <i>6 hours per benefit period</i>
Hearing Loss Screening: Newborns	The USPSTF recommends screening for hearing loss in all newborn infants.	92551 - 92552, 92585 - 92588, V5008	None	Newborns age 0-3 months. <i>1 time per benefit period</i>
Hemoglobinopathies Screening: Newborns	Recommends screening for sickle cell disease in newborns.	83020 - 83021, S3620	None	Newborns age 0-3 months. <i>1 time per benefit period</i>

PREVENTIVE CARE SERVICES GUIDELINES (continued)

SERVICE & DESCRIPTION	USPSTF RECOMMENDATION	CPT/HCPCS CODES	ICD-9-CM CODES	FREQUENCY & CRITERIA
Hepatitis B Screening: Pregnant Women	The USPSTF strongly recommends screening for hepatitis B virus infection in pregnant women at their first prenatal visit.	87340 - 87350, 87515 - 87517	Payable with diagnosis of pregnancy	Pregnant Women. <i>1 time per benefit year</i>
HIV Screening	The USPSTF strongly recommends that clinicians screen for human immunodeficiency virus (HIV) all adolescents and adults at increased risk for HIV infection.	86701 - 86703, 87389 - 87391, 86689, 87534 - 87539, G0432 - G0435, S3645	None	Men and Women age 13 + with an increased risk. <i>1 time per benefit period</i>
Hypothyroidism Screening: Newborns	The USPSTF recommends screening for congenital hypothyroidism in newborns.	84436 - 84439, 84443, S3620	None	Newborns age 0-3 months. <i>1 time per benefit period</i>
Iron Supplementation in Children	The USPSTF recommends routine iron supplementation for asymptomatic children aged 6 to 12 months who are at increased risk for iron deficiency anemia.	Service is included in a preventive care visit. Codes: 99381 - 99429 appropriate to the service	None	Children age 6 – 12 months old.
Obesity Screening and Counseling: Adults	The USPSTF recommends that clinicians screen all adult patients for obesity and offer intensive counseling and behavioral interventions to promote sustained weight loss for obese adults.	97802 - 97804, S9470, G0270 - G0271, G0446 – G0447, 98960 or may be included in a preventive care visit. Codes: 99381 - 99429 appropriate to the service	None for the screening. The counseling is payable with an Obesity diagnosis	Men and Women age 18+. <i>6 hours per benefit period</i>

PREVENTIVE CARE SERVICES GUIDELINES (continued)

SERVICE & DESCRIPTION	USPSTF RECOMMENDATION	CPT/HCPCS CODES	ICD-9-CM CODES	FREQUENCY & CRITERIA
Obesity Screening and Counseling: Children	The USPSTF recommends that clinicians screen children aged 6 years and older for obesity and offer them or refer them to comprehensive, intensive behavioral interventions to promote improvement in weight status.	97802 - 97804, S9470, G0270 - G0271, G0446 – G0447, 98960 or may be included in a preventive care visit. Codes: 99381 - 99429 appropriate to the service	None for the screening. The counseling is payable with an Obesity diagnosis	Children age 6+. 26 hours per benefit period
Osteoporosis Screening: Women	The USPSTF recommends that women aged 65 and older be screened routinely for osteoporosis. The USPSTF recommends that routine screening begin at age 60 for women at increased risk for osteoporotic fractures.	76977, 77078 - 77083, G0130	V17.81, V82.81 or payable with diagnosis related to bone disease	Women age 60 +. 1 time per benefit period
PKU Screening: Newborns	The USPSTF recommends screening for phenylketonuria (PKU) in newborns.	84030, S3620	None	Newborns age 0-3 months. 1 time per benefit period
Rh Incompatibility Screening: First Pregnancy Visit and 24-28 Weeks Gestation	The USPSTF strongly recommends Rh (D) blood typing and antibody testing for all pregnant women during their first visit for pregnancy-related care, and repeating Rh (D) antibody testing for all unsensitized Rh (D)-negative women at 24-28 weeks' gestation, unless the biological father is known to be Rh (D)-negative.	86901, 86850, 86870	Payable with diagnosis of pregnancy	Pregnant Women 24 – 48 weeks gestation and first prenatal visit. 2 per benefit period

PREVENTIVE CARE SERVICES GUIDELINES (continued)

SERVICE & DESCRIPTION	USPSTF RECOMMENDATION	CPT/HCPCS CODES	ICD-9-CM CODES	FREQUENCY & CRITERIA
STIs Counseling	The USPSTF recommends high-intensity behavioral counseling to prevent sexually transmitted infections (STIs) for all sexually active adolescents and for adults at increased risk for STIs.	G0445 or with preventive care visit codes: 99381 - 99429 appropriate to the service	V65.44 - V65.45, V69.2 or V01.6	Men and Women age 13+. <i>3 hours per benefit period</i>
Tobacco Use Counseling: Pregnant Women and Non-Pregnant Adults	The USPSTF recommends that clinicians ask all adults about tobacco use and provide tobacco cessation interventions for those who use tobacco products. Clinicians should also ask all pregnant women about tobacco use and provide augmented, pregnancy-tailored counseling to those who smoke.	G0436 - G0437, S9075, S9453, 99078 or with preventative care visit codes: 99381-99429 appropriate to the service	Payable with diagnosis of 305.1 or V15.82	Men and Women ages 18 +. <i>1 time per benefit period</i>
Syphilis Screening: Non-Pregnant Persons and Pregnant Women	The USPSTF strongly recommends that clinicians screen persons at increased risk for syphilis infection including pregnant women.	86592 - 86593, 86780, 87285.	Payable with diagnosis of pregnancy or with V22.0 - V23.9 or 630-679.14, or V65.44 - V65.45, V69.2 or V01.6	Men and Women ages 18+. <i>1 time per benefit period</i>
Visual Acuity Screening in Children	The USPSTF recommends screening to detect amblyopia, strabismus, and defects in visual acuity in children younger than age 5 years.	92002 - 92014 or S0620 - S0621	V72.0	Children ages 0-5 years. <i>1 time per benefit period</i>

PREVENTIVE CARE SERVICES GUIDELINES (continued)

SERVICE & DESCRIPTION	USPSTF RECOMMENDATION	CPT/HCPCS CODES	ICD-9-CM CODES	FREQUENCY & CRITERIA
TB Testing	USPSTF recognizes the importance of targeted screening for tuberculosis. Per the Centers for Disease Control and Prevention (CDC).	86480, 86580	None	Men and Women ages 18+. <i>1 time per benefit period</i>

These guidelines represent a compilation of recommendations from the United States Preventive Services Task Force (USPSTF). SHPO will make any additions or revisions in accordance with changes in the recommendations of the USPSTF. SHPO reserves the right to amend CPT/HCPCS and ICD-9-CM codes for the Preventive Care Services at its discretion. These additions or revisions will be posted on SHPO's website.

Services listed in the Preventive Care Services Guidelines table are covered with a \$0.00 member cost share when received from a SHPO in-network provider. The guidelines should not replace clinical judgment or the needs of the individual patient.

Publication History

01/01/2012 Effective Date